




## EXHIBIT D



Entrez PubMed Nucleotide Protein Genome Structure PMC Journals B

Search PubMed for

Limits Preview/Index History Clipboard Details

About Entrez

Text Version

Entrez PubMed  
Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-Utilities


PubMed Services  
Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
LinkOut  
Cubby

Related Resources  
Order Documents  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

Privacy Policy

Abstract    Text

☐ 1: Comput Chem. 2001 Jul; 25(4): 411-22. [Related Articles, L](#)

**Amino acid residue environments and predictions of residue ty**

**Kim W, Wilbur WJ.**

National Center for Biotechnology Information, National Library of Medicine  
National Institutes of Health, Bethesda, MD 20894, USA.  
[wonkim@ncbi.nlm.nih.gov](mailto:wonkim@ncbi.nlm.nih.gov)

The determination of a protein's structure from the knowledge of its linear chain is one of the important problems that remains as a bottleneck in interpreting the rapidly increasing repository of genetic sequence data. One approach to this problem that has shown promise and given a measure of success is threading. In this approach contact energies between different amino acids are first determined by statistical methods applied to known structures. These contact energies are then applied to a sequence whose structure is to be determined by threading it through various known structures and determining the total threading energy for each candidate structure. That structure that yields the lowest total energy is then considered the leading candidate among all the structures tested. Additional information is often needed in order to support the results of threading studies, as it is well known in the field that contact potentials used are not sufficiently sensitive to allow definitive conclusions. Here, we investigate the hypothesis that the environment of an amino acid residue realized as all those residues not local to it on the chain but sufficiently close spatially can supply information predictive of the type of residue that is not adequately reflected in the individual contact energies. We present evidence that confirms this hypothesis and suggests a high order cooperativity between the residues that surround a given residue and how they interact with it. We suggest a possible application to threading.

PMID: 11459355 [PubMed - indexed for MEDLINE]

Abstract    Text

[Write to the Help Desk](#)